

L13 ANSWER 4 OF 4 MEDLINE

DUPLICATE 3

AN 95229068 MEDLINE

DN 95229068 PubMed ID: 7713439

TI A large-scale **gene-trap** screen for insertional mutations in developmentally regulated genes in mice.

AU Wurst W; Rossant J; Prideaux V; Kownacka M; Joyner A; Hill D P; Guillemot F; Gasca S; Cado D; Auerbach A; +

CS Samuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto, Ontario, Canada.

NC HD-25334 (NICHD)

SO GENETICS, (1995 Feb) 139 (2) 889-99.

Journal code: 0374636. ISSN: 0016-6731.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199505

ED Entered STN: 19950524

Last Updated on STN: 19950524

Entered Medline: 19950518

571 431.64

AB We have used a **gene-trap** vector and mouse embryonic stem (ES) cells to screen for insertional mutations in genes developmentally regulated at 8.5 days of embryogenesis (dpc). From 38,730 cell lines with vector insertions, 393 **clonal** integrations had disrupted active transcription units, as assayed by beta-galactosidase reporter gene expression. From these lines, 290 clones were recovered and injected into blastocysts to assay for reporter gene expression in 8.5-dpc chimeric mouse embryos. Of these, 279 clones provided a sufficient number of chimeric embryos for analysis. Thirty-six (13%) showed restricted patterns of reporter-gene expression, 88 (32%) showed widespread expression and 155 (55%) failed to show detectable levels of expression. Further analysis showed that approximately one-third of the clones that did not express detectable levels of the reporter gene at 8.5 dpc displayed reporter gene activity at 12.5 dpc. Thus, a large proportion of the genes that are expressed in ES cells are either temporally or spatially regulated during embryogenesis. These results indicate that **gene-trap** mutageneses in embryonic stem cells provide an effective approach for isolating mutations in a large number of developmentally regulated genes.